


# ORTHOTIC TREATMENT: STROKE REHABILITATION



Stroke is the second most common cause of death and the leading cause of disability in Europe. <sup>(1)</sup> Of those that survive an acute stroke, it is estimated that about 40% remain dependant on other people for their daily activities. The treatment of Stroke accounts for approximately 5% of total NHS costs with the societal costs being £8.9 billion. <sup>(3)</sup> Of this total annual direct care was estimated at £4.361 billion. Dropped foot is present in around 20% of patients surviving a stroke. <sup>(2)</sup> In this context orthotic treatment is essential where appropriate given that for every £1 spent on orthotics, the NHS could save up to £4. <sup>(8)</sup>

## Orthotists - members of the stroke team

The orthotist has a key role to play in helping manage contractures, optimise safety, mobility and recovery. Good orthotic intervention can enhance and optimise the effect of other services such as physiotherapy and potentially minimise the need for long term care by preventing secondary complications. <sup>(9,10)</sup>

Joint assessment with other therapists will lead to the most effective integrated treatment to optimise recovery. Using orthoses to stabilise or challenge stability in conjunction with a planned programme with regular review of the orthoses to compliment changes as the patient recovers will ensure best outcome. <sup>(10,11)</sup>

The earlier the orthotist can be involved, the greater help can be offered. The orthotist should be involved each step of the way in the rehabilitation journey from acute to early discharge to review in the chronic stage. <sup>(11)</sup>

## Common orthotic treatment in stroke

Prevention of contractures in the presence of muscle imbalance and abnormal tone.

Optimising alignment in stance as an adjunct to physiotherapy input. This can reduce the number of hands and therapists required in early treatment and ensure consistent repeatable alignment, essential for motor learning. This can both reduce costs by need for fewer staff and optimise quality by ensuring good alignment with every stand <sup>(8,10)</sup>

The orthotist can supplement therapy input by fine tuning the demands versus degree of control provided by the orthoses to match patient needs as recovery progresses. <sup>(10)</sup>

Orthoses can be designed to specifically retrain proximal control as part of a planned rehabilitation programme.

In complex situations Orthotic input can maximise effect of tone management and if necessary functionally accommodate for loss of range. <sup>(11)</sup>

Where the aim of treatment is to have an immediate improvement on walking speed, efficiency or gait pattern or weight bearing during stance, patients should be assessed for suitability for an Ankle Foot Orthosis (AFO) by an orthotist. <sup>(4,5)</sup>

In chronic stroke, the orthotist has the ability to optimise function, allowing for compensations if necessary.

## REFERENCES

- 1) Murray, C.J.L., Lopez, A.D. Global mortality, disability and the contribution of risk factors. Global burden of the disease study. Lancet 1997; 349:1436-42
- 2) Healthcare Improvement Scotland; Update of Evidence note 25: The use of functional electrical stimulation (FES) in adults with dropped foot; Number 46; August.
- 3) Saka, O., McGuire, A., Wolfe, C. 2008 Cost of stroke in the United Kingdom
- 4) Scottish Intercollegiate Guidelines Network Management of patients with stroke: rehabilitation, prevention and management of complications, and discharge planning. NHS Quality Improvement Scotland June 2014.
- 5) Royal college of physicians National clinical guideline for stroke 2012
- 6) Mant, J., Wade, D., Winner, S. (2004) Health care needs assessment. In: Steven et al, editor. Health care needs assessment :the epidemiological based needs assessment reviews, 2nd edition .oxford: Radcliffe Publishing.
- 7) Mohan, K.M., Wolfe, C.D.A., Rudd, A.G., Heuschmann, P.U., Kolominsky-Rabas, P.L., Grieve, A.P. (2011) Risk and cumulative risk of stroke recurrence.
- 8) Boxer, P. and Flynn, T. 2004 Orthotic Pathfinder: a patient focused strategy and proven implementation plan to improve and expand access to orthotic care services and transform the quality of care delivered. Business solutions, London.
- 9) Condie M.E., Campbell J.H., Martina J.D. 2004 Report of a consensus conference on the orthotic management of stroke patients. Copenhagen: ISPO
- 10) Lennon S, Stokes M, 2009 Pocketbook of Neurological Physiotherapy chapter 16 Orthotic management, Elsevier
- 11) Barnes M, Johnson G, 2008, Upper motor neurone syndrome and spasticity: clinical management and neurophysiology. Chapter 6 Orthoses, splints and casts. Cambridge press.
- 12) International Organization for Standardization, 1989. ISO 8549-1:1989: Prosthetics and orthotics -- Vocabulary -- Part 1: General terms for external limb prostheses and external orthoses.

Produced by the British Association of Prosthetists and Orthotists (BAPO)

For further information please contact:

BAPO Secretariat

Sir James Clark Building, Abbey Mill Business Centre, Paisley, PA1 1TJ

Tel: 0141 561 7217

Email: [enquiries@bapo.com](mailto:enquiries@bapo.com) Pais.2)

[www.bapo.com](http://www.bapo.com)